

---

# SELECTED PULMONARY DISORDERS

---

## CHAPTER 32

### Pneumonia

#### KEY TEACHING POINTS

- Many of the classic physical findings of lobar consolidation—diminished chest excursion, dullness, diminished breath sounds, bronchial breath sounds, and egophony—are accurate signs of pneumonia when present. Nonetheless, they appear in only the minority of patients with proven pneumonia; therefore their *absence* does not affect the probability of disease.
- In patients with cough and fever the presence of normal vital signs (i.e., temperature, pulse rate, respiratory rate, and oxygen saturation) decreases probability of pneumonia.
- The Heckerling scoring scheme combines five independent findings of pneumonia (tachycardia, fever, crackles, diminished breath sounds, and *absence* of asthma) and greatly increases the clinician's diagnostic accuracy for pneumonia.
- The CURB-65 score combines five findings to accurately predict the prognosis of patients with pneumonia, information that becomes essential when making decisions about triage of patients.

---

#### I. INTRODUCTION

Like most of the pulmonary examination, the traditional findings of lobar pneumonia were described in 1819 by Laennec, who wrote that clinicians using his newly invented stethoscope could detect acute pneumonia “in every possible case.”<sup>1</sup> According to traditional teachings, the earliest findings of pneumonia are crackles and diminished breath sounds, followed by dullness to percussion, increased tactile fremitus and vocal resonance, and bronchial breath sounds.<sup>2</sup>

## II. CLINICAL SIGNIFICANCE

### A. INDIVIDUAL FINDINGS

**EBM Box 32.1** reviews the findings from more than 6000 patients presenting with acute fever, cough, sputum production, or dyspnea, all of whom underwent chest radiography (the diagnostic standard for pneumonia). The findings increasing probability of pneumonia, in descending order of their likelihood ratios (LRs), are asymmetric chest expansion (LR = 44.1), egophony (LR = 4.1), cachexia (LR = 4), bronchial breath sounds (LR = 3.3), oxygen saturation of less than 95% (LR = 3.1), percussion dullness (LR = 3), respiratory rate higher than 28/minute (LR = 2.7), crackles (LR = 2.3), diminished breath sounds (LR = 2.2), temperature higher than 37.8° C (LR = 2.2), and abnormal mental status (LR = 1.9).

The only finding decreasing the probability of pneumonia was the finding that all vital signs were normal (LR = 0.3). In many studies, wheezing was found more often in patients *without* pneumonia, primarily because the cause of the acute respiratory complaints in these patients was asthma, not pneumonia.<sup>4,5,17,18</sup>

### B. LAENNEC VERSUS MODERN STUDIES

There are three reasons why the studies in **EBM Box 32.1** contradict Laennec's assertion that physical diagnosis is the perfect diagnostic tool: (1) Patients diagnosed with pneumonia today include those with more mild disease than in Laennec's time, when the only available diagnostic standard was postmortem examination (i.e., his conclusions were drawn from patients with only the most severe disease). (2) Many traditional findings appear only after several days of illness, times when the modern clinician, already familiar with the chest radiograph, often examines patients in a more cursory fashion. In contrast, Laennec examined each of his patients diligently day after day, concluding that bronchial breath sounds and bronchophony usually appeared only after 1 to 3 days of hospitalization, and dullness to percussion appeared only after day 4.<sup>1,19</sup> (3) Antimicrobial medications probably alter the course of the physical findings. For example, in the preantibiotic era, fever usually lasted 7 days in patients with lobar pneumonia;<sup>20</sup> nowadays it usually lasts only 3 or 4 days.<sup>21,22</sup>

Even so, many great clinicians of the past tempered Laennec's enthusiasm and taught that auscultation was an imperfect diagnostic tool. Writing only 20 years after Laennec's treatise, Thomas Addison\* stated it was high time "to strip the stethoscope of the extravagant and meretricious pretensions thrust upon it . . . and to state fairly what it will not, as well as what it will do. . . ."<sup>23</sup>

### C. COMBINED FINDINGS

Combining findings improves the accuracy of bedside examination. One of the best models, validated in four different populations<sup>4,15</sup> scores one point for each of the following five findings: (1) temperature higher than 37.8° C, (2) heart rate more than 100/minute, (3) crackles, (4) diminished breath sounds, and (5) *absence* of asthma. **EBM Box 32.1** shows that a score of 4 or 5 argues compellingly for pneumonia (LR = 8.2), whereas a score of 0 or 1 argues *against* pneumonia (LR = 0.3), which in some groups of patients may reduce the probability of pneumonia enough that a chest radiograph becomes unnecessary (e.g., in patients presenting to a community

---

\*Thomas Addison, the discoverer of adrenal insufficiency, was also a recognized master of percussion and auscultation.



## EBM BOX 32.1

## Pneumonia\*

Finding (Reference) <sup>†</sup>	Sensitivity (%)	Specificity (%)	Likelihood Ratio <sup>‡</sup> if Finding Is	
			Present	Absent
<b>General Appearance</b>				
Cachexia <sup>3</sup>	10	97	4.0	NS
Abnormal mental status <sup>4,6</sup>	12-14	92-95	1.9	NS
<b>Vital Signs</b>				
Heart rate >100/min <sup>3-11</sup>	12-65	60-96	1.8	0.8
Temperature >37.8° C <sup>3-14</sup>	16-75	44-95	2.2	0.7
Respiratory rate >28/ min <sup>5-7,13</sup>	7-36	80-99	2.7	0.9
Oxygen saturation <95% <sup>8,13,14</sup>	33-52	80-86	3.1	0.7
All vital signs nor- mal <sup>5,8,11,15,16</sup>	3-38	24-81	0.3	2.2
<b>Lung Findings</b>				
Asymmetric chest expan- sion <sup>3</sup>	5	100	44.1	NS
Chest wall tenderness <sup>12</sup>	5	96	NS	NS
Percussion dullness <sup>3-5,17,18</sup>	4-26	82-99	3.0	NS
Diminished breath sounds <sup>4,5,9,10,17,18</sup>	7-49	73-98	2.2	0.8
Bronchial breath sounds <sup>4</sup>	14	96	3.3	NS
Egophony <sup>3-5</sup>	4-16	96-99	4.1	NS
Crackles <sup>3-6,9,10,12,17,18</sup>	19-67	36-96	2.3	0.8
Wheezing <sup>4-6,9,12,17,18</sup>	10-36	50-86	0.8	NS
<b>Diagnostic Score<sup>4,15</sup></b>				
0 or 1 findings	7-29	33-65	0.3	—
2 or 3 findings	48-55	—	NS	—
4 or 5 findings	38-41	92-97	8.2	—


\*Diagnostic standard: For *pneumonia*, infiltrate on chest radiograph.

<sup>†</sup>Definition of findings: For *all vital signs* normal, temperature <37.8° C, pulse ≤100/min, respirations ≤20,<sup>5,8,11,15,16</sup> and oxygen saturations >95%;<sup>16</sup> for Heckerling *diagnostic score*, the clinician scores one point for each of the following five findings that are present: temperature >37.8° C, heart rate >100/min, crackles, diminished breath sounds, and absence of asthma.

<sup>‡</sup>Likelihood ratio (LR) if finding present = positive LR; LR if finding absent = negative LR.

NS, Not significant.

[Click here to access calculator](#)



office with cough, in whom the probability of pneumonia is 10% or less, a score of 0 or 1 reduces the probability of pneumonia to 3% or less).

D. PNEUMONIA AND PROGNOSIS

In studies of immunocompetent adults hospitalized with community-acquired pneumonia, the 30-day mortality rate is 4% to 15%. Of the individual findings that predict an increased risk of death (EBM Box 32.2), the most compelling ones are hypotension (LR = 7.6) and hypothermia (LR = 3.5).

Several different scoring schemes combine bedside findings to predict mortality in patients with pneumonia. One of the best validated is the **Pneumonia Severity Index**,<sup>51</sup> which unfortunately has the disadvantage of requiring knowledge of 20 different clinical variables, making it difficult to recall and apply at the bedside. A much simpler rule is the CURB-65 score, based on five prognostic variables<sup>†</sup> identified decades ago by the British Thoracic Society:<sup>32</sup> (1) confusion, (2) blood urea nitrogen (BUN) levels greater than 19 mg/dL (>7 mmol/L), (3) respiratory rate of 30 breaths/minute or higher, (4) hypotension (i.e., diastolic blood pressure ≤60 mm Hg or systolic blood pressure ≤90 mm Hg), and (5) age 65 years or older. The presence of three or more of these CURB-65 variables is associated with increased hospital mortality (LR = 2.6 for three findings, LR = 5.9 for four findings, and LR = 11.1 for five findings; see EBM Box 32.2), whereas the absence of all CURB-65 variables is associated with decreased hospital mortality (LR = 0.2 for 0 findings).

The CURB-65 score requires knowledge of the patient's BUN, which may not be immediately available to office-based clinicians. Related scores that omit laboratory values have also been studied, although less extensively so: a CRB-65 score of 0 (i.e., a score of 0 indicates the patient is younger than 65 years and lacks confusion, tachypnea, and hypotension) decreases the probability of mortality (LR = 0.1), and a CRB score of 2 or higher (i.e., two or more of confusion, tachypnea, and hypotension) increases probability of death (LR = 5.0).<sup>26,27,32,33,38,52-54</sup>

<sup>†</sup>CURB-65 is an acronym for Confusion, Urea, Respiratory rate, Blood pressure, and Age ≥65 years.

**EBM BOX 32.2***Pneumonia: Predictors of Hospital Mortality*


Finding* (Reference)	Sensitivity (%)	Specificity (%)	Likelihood Ratio <sup>†</sup> if Finding Is	
			Present	Absent
General Appearance				
Abnormal mental status <sup>24-28</sup>	48-65	70-87	2.7	0.6
Vital Signs				
Heart rate >100/min <sup>24</sup>	45	78	2.1	NS
Systolic blood pressure <90 mm Hg <sup>25,29-31</sup>	11-41	90-99	7.6	0.8
Hypothermia <sup>25,30</sup>	14-43	93	3.5	NS
Respiratory rate >30/min <sup>25,26,32-34</sup>	41-85	63-87	2.1	0.6
Oxygen saturation <90% <sup>35,36</sup>	18-52	75-96	2.8	NS
CURB-65 Prognostic Score <sup>37-50</sup>				
0 findings	0-16	41-92	0.2	—
1 finding	3-38		0.5	—
2 findings	17-51		NS	—
3 findings	13-61		2.6	—
4 findings	4-35		5.9	—
5 findings	1-12	99-100	11.1	—

\*Definition of findings: For hypothermia, body temperature <36.1° C<sup>25</sup> or <37.0° C;<sup>30</sup> for CURB-65 prognostic score, the clinician scores one point for each of the following findings that are present: confusion, BUN > 19 mg/dL, respiratory rate ≥30/min, low blood pressure (either systolic blood pressure ≤90 mm Hg or diastolic blood pressure ≤60 mm Hg), and age ≥65 years.

<sup>†</sup>Likelihood ratio (LR) if finding present = positive LR; LR if finding absent = negative LR.

NS, Not significant.

[Click here to access calculator](#)

**PNEUMONIA: PREDICTORS OF MORTALITY**

## E. HOSPITAL COURSE

Among survivors of pneumonia, abnormalities of the vital signs—fever, tachycardia, tachypnea, and hypotension—usually become normal within 2 to 4 days.<sup>21,22</sup> After this occurs, subsequent clinical deterioration is rare, and fewer than 1% of patients will require subsequent intensive care, coronary care, or telemetry monitoring.<sup>21</sup> If patients are discharged from the hospital before normalization of vital signs, there is an increased risk of readmission and death.<sup>55-57</sup>

*The references for this chapter can be found on [www.expertconsult.com](http://www.expertconsult.com).*

## REFERENCES

1. Laennec RTH. *A Treatise on the Diseases of the Chest*. (facsimile edition by *Classics of Medicine Library*). London: T. and G. Underwood; 1821.
2. Cabot RC. *Physical Diagnosis*. New York, NY: William Wood and Co.; 1926.
3. Diehr P, Wood RW, Bushyhead J, Krueger L, Wolcott B, Tompkins RK. Prediction of pneumonia in outpatients with acute cough—a statistical approach. *J Chron Dis*. 1984;37(3):215–225.
4. Heckerling PS, Tape TG, Wigton RS, et al. Clinical prediction rule for pulmonary infiltrates. *Ann Intern Med*. 1990;113:664–670.
5. Gennis P, Gallagher J, Falvo C, Baker S, Than W. Clinical criteria for the detection of pneumonia in adults: guidelines for ordering chest roentgenograms in the emergency department. *J Emerg Med*. 1989;7:263–268.
6. Mehr DR, Binder EF, Kruse RL, Zweig SC, Madsen RW, D'Agostino RB. Clinical findings associated with radiographic pneumonia in nursing home residents. *J Fam Pract*. 2001;50(11):931–937.
7. Heckerling PS. The need for chest roentgenograms in adults with acute respiratory illness: clinical predictors. *Arch Intern Med*. 1986;146:1321–1324.
8. Nolt BR, Gonzales R, Maselli J, Aagaard E, Camargo CA, Metlay JP. Vital-sign abnormalities as predictors of pneumonia in adults with acute cough illness. *Am J Emerg Med*. 2007;25:631–636.
9. Nakanishi M, Yoshida Y, Takeda N, et al. Significance of the progression of respiratory symptoms for predicting community-acquired pneumonia in general practice. *Respirology*. 2010;15:969–974.
10. van Vugt SF, Broekhuizen BDL, Lammens C, et al. Use of serum C reactive protein and procalcitonin concentrations in addition to symptoms and signs to predict pneumonia in patients presenting to primary care with acute cough: diagnostic study. *Br Med J*. 2013;346:f2540.
11. O'Brien WT, Rohweder DA, Lattin GE, et al. Clinical indicators of radiographic findings in patients with suspected community-acquired pneumonia: who needs a chest x-ray? *J Am Coll Radiol*. 2006;3:703–706.
12. Singal BM, Hedges JR, Radack KL. Decision rules and clinical prediction of pneumonia: evaluation of low-yield criteria. *Ann Emerg Med*. 1989;18(1):13–20.
13. Holm A, Nexoe J, Bistrup LA, et al. Aetiology and prediction of pneumonia in lower respiratory tract infection in primary care. *Br J Gen Pract*. 2007;57:547–554.
14. Kyriacou DN, Yarnold PR, Soltysik RC, et al. Derivation of a triage algorithm for chest radiography of community-acquired pneumonia patients in the emergency department. *Acad Emerg Med*. 2008;15:40–44.
15. Emerman CL, Dawson N, Speroff T, et al. Comparison of physician judgment and decision aids for ordering chest radiographs for pneumonia in outpatients. *Ann Emerg Med*. 1991;20(11):1215–1219.
16. Khalil A, Kelen G, Rothman RE. A simple screening tool for identification of community-acquired pneumonia in an inner city emergency department. *Emerg Med J*. 2007;24:336–338.
17. Melbye H, Straume B, Aasebo U, Dale K. Diagnosis of pneumonia in adults in general practice. *Scand J Prim Health Care*. 1992;10:226–233.
18. Melbye H, Straume B, Aasebo U, Brox J. The diagnosis of adult pneumonia in general practice. *Scand J Prim Health Care*. 1988;6:111–117.
19. Auenbrugger L. *On Percussion of the Chest (1761), Being a Translation of Auenbrugger's Original Treatise Entitled "Inventum novum ex percussione thoracis humani, ut signo abstrusus interni pectoris morbos detegendi."* (facsimile edition by Johns Hopkins Press). Baltimore, MD: The Johns Hopkins Press; 1936.
20. Sahli H. *A Treatise on Diagnostic Methods of Examination*. Philadelphia, PA: W. B. Saunders; 1911.
21. Halm EA, Fine MJ, Marrie TJ, et al. Time to clinical stability in patients hospitalized with community-acquired pneumonia: implications for practice guidelines. *J Am Med Assoc*. 1998;279:1452–1457.
22. Menendez R, Torres A, de Castro FR, et al. Reaching stability in community-acquired pneumonia: the effects of the severity of disease, treatment, and the characteristics of patients. *Clin Infect Dis*. 2004;39:1783–1790.

23. Addison T. The difficulties and fallacies attending physical diagnosis of diseases of the chest. In: Wilks D, ed. *A Collection of the Published Writings of the Late Thomas Addison*. (facsimile edition by *Classics of Medicine Library*). London: The New Sydenham Society; 1846:242.
24. Starczewski AR, Allen SC, Vargas E, Lye M. Clinical prognostic indices of fatality in elderly patients admitted to hospital with acute pneumonia. *Age Ageing*. 1988;17:181–186.
25. Conte HA, Chen YT, Mehal W, Scinto JD, Qualiarello VJ. A prognostic rule for elderly patients admitted with community-acquired pneumonia. *Am J Med*. 1999;106:20–28.
26. Ewig S, Kleinfeld T, Seifert K, Shafer H, Goke N. Comparative validation of prognostic rules for community-acquired pneumonia in an elderly population. *Eur Respir J*. 1999;14:370–375.
27. Neill AM, Martin IR, Weir R, et al. Community acquired pneumonia: aetiology and usefulness of severity criteria on admission. *Thorax*. 1996;51:1010–1016.
28. Myint PK, Kamath AV, Vowler SL, Maisey DN, Harrison BDW. Severity assessment criteria recommended by the British Thoracic Society (BTS) for community-acquired pneumonia (CAP) and older patients: should SOAR criteria be used in older people? A compilation study of two prospective cohorts. *Age Ageing*. 2006;35:286–291.
29. Fedullo AJ, Swinburne AJ. Relationship of patient age to clinical features and outcome for in-hospital treatment of pneumonia. *J Gerontol*. 1985;40:29–33.
30. Allen SC. Lobar pneumonia in Northern Zambia: clinical study of 502 adult patients. *Thorax*. 1984;39:612–616.
31. Chalmers JD, Singanayagam A, Hill AT. Systolic blood pressure is superior to other haemodynamic predictors of outcome in community acquired pneumonia. *Thorax*. 2008;63:698–701.
32. Research Committee of the British Thoracic Society. Community-acquired pneumonia in adults in British hospitals in 1982–1983: a survey of aetiology, mortality, prognostic factors and outcome. *Q J Med*. 1987;62(239):195–220.
33. Farr BM, Sloman AJ, Fisch MJ. Predicting death in patients hospitalized for community-acquired pneumonia. *Ann Intern Med*. 1991;115:428–436.
34. Brancati FL, Chow JW, Wagener MM, Vacarello SJ, Yu VL. Is pneumonia really the old man's friend? Two-year prognosis after community-acquired pneumonia. *Lancet*. 1993;342:30–33.
35. Bewick T, Greenwood S, Lim WS. What is the role of pulse oximetry in the assessment of patients with community-acquired pneumonia in primary care? *Prim Care Respir J*. 2010;19:378–382.
36. Majumdar SR, Eurich DT, Gamble JM, Senthilselvan A, Marrie TJ. Oxygen saturations less than 92% are associated with major adverse events in outpatients with pneumonia: a population-based cohort study. *Clin Infect Dis*. 2011;52:325–331.
37. Aujesky D, Auble TE, Yealy DM, et al. Prospective comparison of three validated prediction rules for prognosis in community-acquired pneumonia. *Am J Med*. 2005;118:384–392.
38. Capelastegui A, Espana PP, Quintana JM, et al. Validation of a predictive rule for the management of community-acquired pneumonia. *Eur Resp J*. 2006;27:151–157.
39. Lim WS, Macfarlane JT, Boswell TCJ, et al. Study of community acquired pneumonia aetiology (SCAPA) in adults admitted to hospital: implications for management guidelines. *Thorax*. 2001;56:296–301.
40. Ananda-Rajah MR, Charles GGP, Melvani S, Burrell LL, Johnson PDR, Grayson L. Comparing the pneumonia severity index with CURB-65 in patients admitted with community acquired pneumonia. *Scand J Infect Dis*. 2007;40:293–300.
41. Espana PP, Capelastegui A, Quintana JM, et al. Validation and comparison of SCAP as a predictive score for identifying low-risk patients in community-acquired pneumonia. *J Infect*. 2010;60:106–113.
42. Man SY, Lee N, Ip M, Antonio GE, et al. Prospective comparison of three predictive rules for assessing severity of community-acquired pneumonia in Hong Kong. *Thorax*. 2007;62:348–353.
43. Menéndez R, Martínez R, Reyes S, et al. Biomarkers improve mortality prediction by prognostic scales in community-acquired pneumonia. *Thorax*. 2009;64:587–591.
44. Schuetz P, Koller M, Christ-Crain M, et al. Predicting mortality with pneumonia severity scores: importance of model recalibration to local settings. *Epidemiol Infect*. 2008;136:1628–1637.



45. Zuberi FF, Khan JA. Prospective comparison of prediction rules of mortality risk for CAP in a developing country. *Int J Tuberc Lung Dis.* 2008;12(4):447–452.
46. Kim HI, Kim SW, Chang HH, et al. Mortality of community-acquired pneumonia in Korea: assessed with the pneumonia severity index and the CURB-65 score. *J Kor Med Sci.* 2013;28:1276–1282.
47. Abisheganaden J, Ding YY, Chong WF, Heng BH, Lim TK. Predicting mortality among older adults hospitalized for community-acquired pneumonia: an enhanced confusion, urea, respiratory rate and blood pressure score compared with pneumonia severity index. *Respirology.* 2012;17:969–975.
48. Guo Q, Li HY, Zhou YP, et al. CURB-65 score predicted mortality in community-acquired pneumonia better than IDSA/ATS minor criteria in a low-mortality-rate setting. *Eur J Clin Microbiol Infect Dis.* 2012;31:3281–3286.
49. Mbata GC, Chukwuka CJ, Onyedum CC, Onwubere BJ. The CURB-65 scoring system in severity assessment of Eastern Nigerian patients with community-acquired pneumonia: a prospective observational study. *Prim Care Respir J.* 2013;22:175–180.
50. Jeong BH, Koh WJ, Yoo H, et al. Performances of prognostic scoring systems in patients with healthcare-associated pneumonia. *Clin Infect Dis.* 2013;56:625–632.
51. Fine MJ, Auble TE, Yealy DM, et al. A prediction rule to identify low-risk patients with community-acquired pneumonia. *N Engl J Med.* 1997;336:243–250.
52. Bauer TT, Ewig S, Marre R, Suttrop N, Welte T. CRB-65 predicts death from community-acquired pneumonia. *J Intern Med.* 2006;260:93–101.
53. Ewig S, de Roux A, Bauer T, et al. Validation of predictive rules and indices of severity for community acquired pneumonia. *Thorax.* 2004;59:421–427.
54. Lim WS, van der Eerden MM, Laing R, et al. Defining community acquired pneumonia severity on presentation to hospital: an international derivation and validation study. *Thorax.* 2003;58:377–382.
55. Halm EA, Fine MJ, Kapoor WN, Singer DE, Marrie TJ, Siu AL. Instability on hospital discharge and the risk of adverse outcomes in patients with pneumonia. *Arch Intern Med.* 2002;162:1278–1284.
56. Dagan E, Novack V, Porath A. Adverse outcomes in patients with community acquired pneumonia discharged with clinical instability from Internal Medicine Department. *Scand J Infect Dis.* 2006;38:860–866.
57. Capelastegui A, Yandiola PPE, Quintana JM, et al. Predictors of short-term rehospitalization following discharge of patients hospitalized with community-acquired pneumonia. *Chest.* 2009;136:1079–1085.